

# Agricultural Weather Study Interim Report to the Minnesota Department of Commerce

June 2022



UNIVERSITY  
OF MINNESOTA  
**Driven to Discover<sup>SM</sup>**

**Prepared by:**

Dr. Heidi Roop, Dr. Tracy Twine, and Amanda Farris  
University of Minnesota, Department of Soil, Water, and Climate  
University of Minnesota Climate Adaptation Partnership

Per the requirements set forth in Minnesota Statute 3.197, the cost to prepare this report was \$830.



# Executive Summary

The purpose of this report is to provide an overview of the work completed on the Agricultural Weather Study through June 2022. The Agricultural Weather Study is advancing towards the goals of developing dynamically downscaled climate projections for the state of Minnesota and creating online resources, an interactive data tool, and dedicated training opportunities to ensure the usefulness and usability of these data to inform a range of climate-related decision-making across the state. We have made significant progress since the project was launched in October 2021 and all current project tasks are in progress or on track.

## Project Overview

Assumptions about climate are embedded in how we design and manage many of the socio-economic systems and resources we rely on every day. Observed climate data and climate models are used to inform how we design our infrastructure, when to plant our crops, how we design railways and bridges, how we assess risk of infectious diseases, etc. But the climate of the past is no longer the best predictor of future climate. To better understand future climate risks, we use future climate models. The Minnesota State Legislature supported the University of Minnesota to generate future climate projections, as well as associated resources to support the use and application of these data across Minnesota's diverse sectors from agriculture and forestry to water resources management and the built environment.

Differentiating the impacts of climate change at a fine spatial scale is particularly important in Minnesota, where we see large south-to-north gradients in warming, have long-duration snow cover, large and small lakes, and other complexities in our climate. In order to provide reasonable climate projections over Minnesota on the regional scale, we need information that captures these complexities. This project will produce climate projections at approximately 3-mile resolution for the entire state using regional climate downscaling techniques (See Task 1). These data will be made available to the public through an interactive web tool and will include projections for three future climate scenarios for three time-steps out to the year 2100.

Results from a recent statewide survey ([Clark et al., 2021](#)) and a subgroup of the State's [Resiliency and Adaptation Action Team](#), point to broad, cross-sectoral demand for these projections and increased support for using this information in practice. Over 80% of respondents to the statewide survey felt that fine scale climate projections were important for carrying out their work. In addition to high demand from State agency staff, researchers, nonprofits and the private sector have all articulated needs and applications for this critical, forward-looking climate information. Anticipated applications span everything from agricultural decision-making to infrastructure design and community engagement. Leveraging the University's research and Extension capacity, this project was designed to both generate the model output and assist in the use of this needed information in risk management and related decision-making.

## Highlights

The report is structured primarily around the tasks in the project contract. Highlights of the project to-date are summarized below.

## 1) Modeling and Data Visualization

- The computer modeling using the Minnesota Supercomputing Institute (MSI) at the University of Minnesota is underway. We are currently running six global climate models and a 1D lake model out to the year 2100 under three different future climate scenarios. We are using the Weather Research and Forecasting Model (WRF) to generate the dynamically downscaled climate projections at approximately 3-mile resolution (4 km).
- The University of Minnesota's U-Spatial team has assigned developers and data visualization experts to begin designing the interactive data tool.
- In collaboration with U-Spatial and the University of Minnesota Office of Information Technology, we are conducting focus groups with data users and using affinity diagramming to capture user stories to inform the design and anticipated uses of the interactive data tool.

## 2) Capacity and Training

- A new Climate Resilience Researcher was hired in February 2022 to help support community engagement and communications material development and to assist with training development and design. This researcher is actively working to expand partnerships, develop training materials, coordinate a project evaluation design, and develop a workflow for the train-the trainer events that will be implemented in 2023.
- A new Climate Resilience Extension Educator, who will be focused on agriculture, was hired in June and will be working with the Agriculture Weather Study project team. This educator will help agricultural producers to understand and use this, and other, climate information in their work and decision-making. This Extension Educator is supported by both this project and the Minnesota Corn Growers Association. This support is enabling our team to develop dedicated capacity, resources, and training to support a new climate-smart agriculture extension program. The Minnesota Corn Growers Association and Minnesota Department of Agriculture both had representatives sit on the search committee for this position.
- Widespread interest in these data and use across sectors, including engineering and architecture, has resulted in a dedicated summer internship opportunity for a University of Minnesota architecture student to work with the project team and experts at the Minnesota-founded architecture and engineering firm Hammel, Green and Abrahamson (HGA). This internship is currently underway and will include a survey of climate data user and data needs across the architecture, engineering, and urban planning disciplines. This work will facilitate the translation and use of this project's output into architecture, engineering and design.

## 3) Communication, Engagement and Future Planning

- We are sharing this project with a range of audiences across the state and Midwest region. To-date this has included several conversations with public and private sector end users throughout Minnesota and with other end users and experts in the Midwest, and nationally. This included a recent presentation to colleagues with the National Weather Service, National Oceanic and Atmospheric Administration (NOAA), US Department of Agriculture and more at the 2022 Climate Prediction Applications Science Workshop (CPASW). We have also been invited to

share this work and the broader programming of the University of Minnesota Climate Adaptation Partnership (MCAP) at the upcoming National Adaptation Forum in October 2022.

- We are also sharing this research internally at the University of Minnesota through venues like the MSI Research Exhibition 2022 and are actively working to identify funding opportunities to further support the use and integration of these data in additional modeling efforts, such as hydrologic and ecosystem modeling, through other grant and research proposals.

## Next Steps and Challenges

During the coming months, the project team will continue to make progress to meet the goals of this project. At present, we do not foresee any major challenges as we have all of the project tasks started or in the planning phases. The computer modeling is well underway and we are completing preliminary testing of the weather models and initializing routines for all of the model runs. We anticipate with current progress that we will be able to complete all model runs to develop the full dataset of downscaled climate projections, with initial data anticipated in November 2022.

As the climate modeling work is performed, we are advancing a range of next steps, including:

- Continuing to engage stakeholders to better understand climate data information needs and to inform development of the interactive web tool and train-the-trainer events.
- Applying methodologies from software engineering, including affinity diagramming and user stories, in conjunction with focus groups, interviews and surveys to capture stakeholder input to inform tool design.
- Developing educational resources and train-the-trainer opportunities that will be aimed at increasing the understanding of the climate projections and their applications in a diversity of decision contexts. These opportunities will include training in the use of the interactive web tool.
- Developing an evaluation plan informed by the rapidly evolving literature on climate services and knowledge co-production with data users, and geared toward continuous improvement of data usability and usefulness over time.

As we look towards the anticipated project completion in December, 2023, we see two potential long-term challenges. The first includes securing sufficient resources for hosting and maintaining the data and interactive tool long-term. Even after the full release of the tool, we anticipate ongoing maintenance and support will be required to ensure usability and function of the interactive web tool. There are also ongoing costs related to the data storage and for hosting the tool online. We are actively in dialogue about long-term solutions to this challenge. A second challenge is related to ensuring the ability to provide the ongoing support and requests for technical services that will stem from the generation of these data after the project is complete. Demand for such support is growing and we are actively seeking pathways to ensure continuity of the positions and staff capacity created by this project so the team is positioned to be able to provide end-users with the needed technical support required to ensure broad use and application of the climate projections across the entire State.

# Tasks and Deliverables

## Task 1. Computer Modeling

This task requires using resources at the Minnesota Supercomputing Institute to analyze high-performing weather model projections under at least two greenhouse gas emissions scenarios and develop a series of projections of temperature, precipitation, snow cover, and other climate parameters out to the year 2100 for the state of Minnesota at a scale as small as three square miles (~4 km).

### Subtask 1.1. Select model scenarios for analysis

Status: We are currently running six global climate models from the Coupled Model Intercomparison Project 6 ([CMIP6](#); Eyring et al., 2016) for this project. These global scenarios will be used in the downscaling of the climate projections to the state of Minnesota out to the year 2100 at approximately 3 mile resolution. This downscaling will be completed using the Weather Research and Forecasting Model (WRF). We have selected three different future climate 'pathways' intended to span a range of possible future outcomes, including ambitious adoption of adaptation and mitigation measures globally (less planetary warming) to continued development of fossil fuels (more planetary warming). We will be using Shared Socioeconomic Pathways (SSPs) as the future climate scenarios following the CMIP6 modeling process and the Intergovernmental Panel on Climate Change (IPCC). This work will update the downscaling of the previous generation of global climate model projections for Minnesota (Liess et al., 2022; Appendix A).

<i>Subtask 1.1 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Select model scenarios	1/1/2023	Complete

### Subtask 1.2. Identify and delineate projection parameters

Status: In parallel with testing our simulations on the Supercomputer, we are consulting with other regional climate data experts and climate data end users at a range of organizations, businesses and state agencies to understand use cases, data needs and priority variables for inclusion in the interactive tool. These discussions include identification of the climate variables of most relevance to stakeholders as well as discussions on model characteristics including selection of a lake model to represent lake surface temperatures and temperatures within lakes.

This engagement work is ongoing and will continue throughout 2022. Examples of discussions we have had to date include staff at entities such as the Minnesota Department of Natural Resources, Minnesota Pollution Control Agency, Minnesota Department of Health, Metropolitan Council, Hennepin County, HGA and Stantec Engineering, the 1854 Treaty Authority. The list of engagement and conversations is actively growing as we have new project members joining the team, increased awareness of the data and as we scope out gaps and needs for additional engagement and dialogue. We are also engaging experts from other states who have led the development of similar data projects and tools to help incorporate lessons learned throughout this project.

<i>Subtask 1.2 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Identify projection parameters	1/1/2023	On track

### Subtask 1.3. Perform projections

Status: Simulations are currently underway at the Minnesota Supercomputing Institute. We are pre-processing the CMIP6 models, initializing the WRF model to ingest these data, and testing the model to correctly perform across the state. We are increasing the horizontal resolution from our previous project (Liess et al. 2022) by a factor of 6 per area from 10km x 10km (~6 miles) to 4km x 4km (~3 miles). By the end of the project we will have improved projections through the 21st century with the most recent scenarios of human greenhouse gas emissions and corresponding climate. For more information on the previous work using the Coupled Model Intercomparison Project 5 model products and a similar downscaling technique, see Appendix A.

<i>Subtask 1.3 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Model analysis and series of year 2100 projections	1/1/2023	On track

## Task 2. Interactive Tool Development

This task will lead to the development of an interactive data tool to facilitate access and use of the model projection data. The tool will be hosted by the University of Minnesota Climate Adaptation Partnership on the following website: [climate.umn.edu](https://climate.umn.edu).

### Subtask 2.1. Outline goals for user interface and experience

Status: We are partnering with U-Spatial and the University of Minnesota Office of Information Technology (OIT) to conduct focus groups with key data users and decision leaders across a broad range of sectors to identify data needs and guide development of the interactive data tool. We have developed a work plan utilizing affinity diagramming and user stories as methods for capturing and applying input from these focus groups to optimize user experience with the tool and usability of data provided for targeted resilience actions. Surveys and interviews of engineering and architecture professionals conducted by the graduate student intern co-mentored with HGA will provide additional input specific to the building sector.

<i>Subtask 2.1 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Outline goals for UI/UX	12/31/2023	On track

### Subtask 2.2. Develop interface components

Status: We are conducting a review of existing online climate data tools to identify common interface components. Further input on components to include in the interactive data tool will be derived from focus group discussions and results from affinity diagramming and crafting of user stories, described in Subtask 2.1. We are also nearing the completion of a range of preliminary communication materials related to climate change projections and climate modeling. These resources are intended to help

end-users of the data have access to easy-to-understand resources about the source of the data, the downscaling methodology and climate projections basics.

<i>Subtask 2.2 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Develop interface components	12/31/2023	On track

#### Subtask 2.3. Perform preliminary testing on user interface

Status: We have not yet started any user interface testing as the tool is not yet built while we wait for the climate modeling task to be completed. Tool design and user engagement are underway to inform the eventual user testing process (see Task 2.1).

<i>Subtask 2.3 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Functional interactive data tool	12/31/2023	On track

### Task 3. Training

Task 3 entails designing and holding at least two train-the-trainer events to increase access to, and understanding of the data, and how to use and apply them in different contexts. This will include providing written materials and communication resources about the data and their use for others to access outside of these training opportunities.

#### Subtask 3.1. Identify and publicize training opportunities to potential stakeholders

Status: Input on training needs, priority user groups, and potential training opportunities will be derived from focus group discussions and results from affinity diagramming and crafting of user stories, described in Subtask 2.1. Surveys and interviews of engineering and architecture professionals conducted by the graduate student intern co-mentored with HGA partners will provide additional input on training needs and opportunities specific to the building sector.

<i>Subtask 3.1 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Identify and publicize training opportunities	10/31/2023	On track

#### Subtask 3.2. Schedule and host training events

Status: We are actively scoping opportunities to conduct at least two train-the-trainer events in mid-2023 and also anticipate leveraging other opportunities including the University of Minnesota Climate Adaptation Partnership's monthly webinar series and anticipated new climate-ready for agriculture and other related Extension programming to engage additional end-users and public audiences in the availability and use of these data.



<i>Subtask 3.2 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Schedule and host training events	10/31./2023	On track

Subtask 3.3. Solicit and compile feedback from attendees, address questions and incorporate results into reporting and data tool refinements

Status: Encompassing our UX/UI processes described above we are planning a developmental evaluation that will start with identifying a group of research and experience-based principles for design effectiveness to guide the creation of data tool prototypes. These principles will be refined initially through UX/UI processes, and then through evaluative methods like surveys at the point-of-download or brief interviews with users. Over the course of the project, we expect these principles will resolve to guide effective design for our range of different user groups and sectors.

<i>Subtask 3.3 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Functional interactive data tool	10/31./2023	On track

## Task 4. Project Reporting

This task relates to project reporting.

Subtask 4.1. Provide an interim report highlighting project progress and results to date.

<i>Subtask 4.1 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Interim report	6/30/2022	Complete; this report

Subtask 4.2. Produce a written final report that includes sufficient detail for technical readers and a clearly written summary for nontechnical readers.

<i>Subtask 4.2 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Written final report	12/31/2023	On track

Subtask 4.3. Final reports, any mid-project status reports, and renewable development account financial reports will be posted online on a public website designated by the commissioner of commerce and will adhere to State accessibility standards.

<i>Subtask 4.3 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Publicly available report materials	12/31/2023	On track

## Task 5. Project updates and invoices

This task entails regular status updates and supplying project invoices. Regular updates and required invoicing are occurring in accordance with the project contract and requests from the Project Manager at the Minnesota Department of Commerce.

Subtask 5.1. Conduct phone conferences as needed with the State's Authorized Representative to apprise him/her of progress, accomplishments and issues encountered.

<i>Subtask 5.1 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Regular phone conferences with State's Authorized Representative	Ongoing	On track

Subtask 5.2. Schedule project update meetings as necessary to inform the State's Authorized Representative of deviations to the project schedule, the need to modify the scope of the project or at the request of the State's Authorized Representative to discuss any item related to the project's progress.

<i>Subtask 5.2 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Update State's Authorized Representative on deviations from schedule or scope	Ongoing	On track

### Subtask 5.3. Quarterly Reporting and Status Updates

5.3.1 Status updates to the State for the preceding period's work detailing progress made toward completing individual project tasks as well as any deviations from the project schedule.

5.3.2 Invoices and supporting documentation to the State for the preceding period's work completed within the project scope; and

5.3.3 Budget overview for the preceding period's expenses and expenses to date

<i>Subtask 5.3 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
5.3.1 Status updates	Ongoing	On track
5.3.2 Invoices	Ongoing	On track
5.3.3 Budget review with State's Authorized Representative	Ongoing	On track

Subtask 5.4. Upon project completion, submit the final invoice with supporting documentation.

<i>Subtask 5.4 Deliverables</i>	<i>Due Date</i>	<i>Status</i>
Final invoice	12/31/2023	On track

## Appendix A

High-Resolution Climate Projections Over Minnesota for the 21st Century Liess et al., 2022 publication

# Earth and Space Science



## RESEARCH ARTICLE

10.1029/2021EA001893

### Key Points:

- Over northern and central Minnesota, winters and summers may be up to 6 and 4°C warmer, respectively, at the end of the 21st century
- Spring precipitation may increase by more than 1 mm per day over northern Minnesota
- Snow depth may decrease by more than 12 cm. Number of snow days per year may decrease by up to 55

### Supporting Information:

Supporting Information may be found in the online version of this article.

### Correspondence to:

S. Liess,  
[liess@umn.edu](mailto:liess@umn.edu)

### Citation:

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-resolution climate projections over Minnesota for the 21st century. *Earth and Space Science*, 9, e2021EA001893. <https://doi.org/10.1029/2021EA001893>

Received 25 JUN 2021

Accepted 2 FEB 2022

### Author Contributions:

**Conceptualization:** Stefan Liess, Tracy E. Twine, Peter K. Snyder

**Data curation:** Stefan Liess, Gabriel Konar-Steenberg

**Formal analysis:** Stefan Liess, Tracy E. Twine, Gabriel Konar-Steenberg

**Funding acquisition:** Tracy E. Twine, Peter K. Snyder, William D. Hutchison, Bonnie L. Keeler, Kate A. Brauman

**Investigation:** Stefan Liess, Tracy E. Twine, Peter K. Snyder, Gabriel Konar-Steenberg

© 2022 The Authors. Earth and Space Science published by Wiley Periodicals LLC on behalf of American Geophysical Union.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

## High-Resolution Climate Projections Over Minnesota for the 21st Century

Stefan Liess<sup>1</sup> , Tracy E. Twine<sup>1</sup> , Peter K. Snyder<sup>1</sup> , William D. Hutchison<sup>2</sup> , Gabriel Konar-Steenberg<sup>1</sup> , Bonnie L. Keeler<sup>3</sup> , and Kate A. Brauman<sup>4,5</sup> 

<sup>1</sup>Department of Soil, Water, and Climate, University of Minnesota, St. Paul, MN, USA, <sup>2</sup>Department of Entomology, University of Minnesota, St. Paul, MN, USA, <sup>3</sup>Humphrey School of Public Affairs, University of Minnesota, St. Paul, MN, USA, <sup>4</sup>Institute on the Environment, University of Minnesota, St. Paul, MN, USA, <sup>5</sup>Global Water Security Center, The University of Alabama, Tuscaloosa, AL, USA

**Abstract** Minnesota is the state with the strongest winter warming in the contiguous United States. We performed regional climate projections at 10 km horizontal resolution using the Weather Research Forecasting model forced with eight CMIP5 GCMs. The selected GCMs have previously been found to be in relatively good agreement with observations over Minnesota compared to other members of the CMIP5 model ensemble. Our projections suggest ongoing warming in all seasons, especially in winter, as well as shallower snow depth and fewer days with snow cover. We expect significant increases in spring and early summer heavy precipitation events. Our comparisons between different time slices and two different emission scenarios indicate a climate for the state of Minnesota near the end of the 21st century that is significantly different from what has been observed by the end of the 20th century. Winters and summers are expected to be up to 6 and 4°C warmer, respectively, over northern and central Minnesota, and spring precipitation may increase by more than 1 mm d<sup>-1</sup> over northern Minnesota. Especially over the central part of the state, winter snow depth is projected to decrease by more than 12 cm, and the number of days per year with snow depth of more than 2.54 cm (one inch) is expected to decrease by up to 55.

**Plain Language Summary** Minnesota is the state with the strongest winter warming in the contiguous United States. We performed regional projections of the climate across Minnesota for the middle and end of the 21st century. We selected the results from eight recent global climate model projections to calculate climate data over an area of 10 km by 10 km with a regional climate model. Our results indicate that the future climate for the state of Minnesota is likely to be significantly different from what has been observed near the end of the 20th century. Over northern and central Minnesota, winters and summers are expected to be up to 6 and 4°C warmer, respectively, near the end of the 21st century. Spring precipitation may increase by more than 1 mm d<sup>-1</sup> over northern Minnesota. Over the central part of the state, winter snow depth is suggested to decrease by more than 12 cm. The number of days per year with snow depth of more than 2.54 cm (one inch) is expected to decrease by up to 55. These results are expected to influence regional decision-making related to agriculture, infrastructure, water resources, and other sectors.

## 1. Introduction

For almost a century, surface warming has had its largest amplitude in the higher latitudes of the Northern Hemisphere (Callendar, 1938). Over the last several decades, the Arctic region has warmed between 0.14°C per decade (Bekryaev et al., 2010) and 0.17°C per decade (Polyakov et al., 2002), which is more than twice the rate of the rest of the planet. With anthropogenic climate change, it may warm an additional 4–8°C by the end of this century (Harvey et al., 2015; IPCC, 2013, 2021). The U.S. state of Minnesota is strongly affected by this Arctic warming, particularly during winter, when the influence of Arctic air is most dominant (Wang et al., 2017) and when reductions in snow cover lower the surface albedo (Shi et al., 2013). Minnesota's winter warming is the strongest among the 48 contiguous United States (NCEI, 2021). Future projections indicate ongoing warming as well as significant increases in spring and early summer heavy precipitation events over the north central United States by the end of this century (Harding & Snyder, 2014).

Despite the clear signals of increased temperature and precipitation in models of future climate, the large-scale outputs of general circulation models (GCM) are difficult to integrate into regional, state, and local planning

**Methodology:** Stefan Liess, Tracy E. Twine, Peter K. Snyder, William D. Hutchison, Bonnie L. Keeler  
**Project Administration:** Tracy E. Twine, Peter K. Snyder, William D. Hutchison, Bonnie L. Keeler  
**Resources:** Stefan Liess, Tracy E. Twine, Peter K. Snyder, William D. Hutchison, Bonnie L. Keeler, Kate A. Brauman  
**Software:** Stefan Liess, Tracy E. Twine, Peter K. Snyder, Gabriel Konar-Steenberg  
**Supervision:** Tracy E. Twine, Peter K. Snyder, William D. Hutchison, Bonnie L. Keeler  
**Validation:** Stefan Liess, Tracy E. Twine, Gabriel Konar-Steenberg  
**Visualization:** Stefan Liess, Gabriel Konar-Steenberg  
**Writing – original draft:** Stefan Liess  
**Writing – review & editing:** Stefan Liess, Tracy E. Twine, Peter K. Snyder, Gabriel Konar-Steenberg, Kate A. Brauman

where climate information is required by decision makers over smaller areas, such as individual watersheds and counties. GCMs typically have resolutions coarser than 100 km, which is insufficient for these applications (Boé et al., 2007; Zorita & von Storch, 1999). Differentiating the impacts of climate change at finer spatial scales is particularly important and challenging in Minnesota, where many days of snow cover and many small-scale open water sources, such as lakes and rivers all contribute to variations on the ground that are typically not addressed by GCMs. In order to provide reasonable climate projections over Minnesota on the regional scale, we dynamically downscaled GCM projections from an eight-model ensemble to a higher spatial resolution (~10 km) by nesting a finer scale regional climate model (RCM).

An early review of dynamical downscaling efforts (Giorgi & Mearns, 1991) describes simulations as high as  $0.5^\circ \times 0.5^\circ$  horizontal resolution, which has later been improved to a range of 25–50 km in the comprehensive Coordinated Regional Downscaling Experiment (CORDEX; Giorgi et al., 2015), especially the North American branch (NA-CORDEX; McGinnis & Mearns, 2021), which is preceded by the North American Regional Climate Change Assessment Program (NARCCAP; Mearns et al., 2009). However, these simulations have an insufficient horizontal resolution to resolve the small-scale open water sources in Minnesota.

Other high-resolution studies like Liu et al. (2017) at a 4-km horizontal resolution typically do not provide a multimodel ensemble to address the uncertainty in future climate projections. Ensemble downscaling simulations are considered crucial for providing a better estimate of future climate change and an uncertainty range (Xu et al., 2018). The downscaling experiment by Ashfaq et al. (2016) is probably closest to the present study. It uses 11 GCMs, including seven of the eight in the present study, for dynamical downscaling over the contiguous United States at an 18-km horizontal resolution, but only addresses one midcentury RCP8.5 scenario.

## 2. Methods

We use a dynamical downscaling approach based on nesting GCM input data with the Weather Research and Forecasting (WRF) RCM (Skamarock et al., 2008) coupled to the Community Land Model (CLM; Dai et al., 2003) with a dynamic crop module. This model version, also known as WRF-CLM4crop, has previously been described by Harding et al. (2016) and Lu et al. (2015). Vegetation, soil, and other land surface parameters for WRF-CLM4crop are taken from the annual cycle of the Moderate-resolution Imaging Spectroradiometer (MODIS) satellite product at a 30-s horizontal resolution (Zhang et al., 2006) and kept consistent between all simulations. Atmospheric boundary conditions include 6-hourly prognostic variables such as temperature, wind speed and direction, specific humidity, and geopotential height from the Coupled Model Intercomparison Project 5 (CMIP5) GCM archive (Taylor et al., 2012) at the Earth System Grid (Williams et al., 2009).

From more than 40 available GCMs, we selected eight that provide all necessary prognostic variables and show reliable large-scale results over the Midwestern United States (Table 1; Harding et al., 2013). We adjusted the model selection in Table 4 of Harding et al. (2013) to only include GCMs that produce reasonable downscaling results during all four seasons, since Harding et al. (2013) only studied the summer season. Therefore, we needed to exclude MIROC4h and the ACCESS models after a short test simulation with our WRF downscaling setup. CCSM4 is not listed in Table 4 of Harding et al. (2013) because 6-hourly input data were not available at that time. However, it is ranked highly in their Figure 5, so we included it in this study.

Our downscaling approach is based on one-way nesting between the global and regional domains and two-way nesting between the two regional domains, so our RCM results cannot modify the global domain. Our regional domains comprise an outer nest over a large portion of North America at a 50-km grid-cell horizontal resolution and an inner nest over Minnesota and adjacent areas at a 10-km grid-cell horizontal resolution (Figure 1). These two nests are connected via two-way nesting and thus influence one another. The southern end of the outer nest is extended to the Gulf of Mexico in order to capture the Great Plains Low Level Jet (GPLLJ), which transports much-needed moisture from the Gulf of Mexico into the Central United States, especially during the warmer seasons (Zhou et al., 2021). However, the extent in other directions is limited by available computing resources, and therefore other storms such as those that develop in the lee of the Rocky Mountains may be represented only by the GCM input data instead of the regional WRF simulations. The inner nest in Figure 1 includes all lake points as represented in our WRF version.

**Table 1**  
*List of GCMs for Boundary Conditions*


Model	Institution	Resolution [°]	Reference
bcc-csm1-1	BCC (China)	1 x 1.33	(Wu et al., 2010)
CCSM4	NCAR (USA)	0.9 x 1.25	(Gent et al., 2011)
CMCC-CM	CMCC (Italy)	0.75 x 0.75	(Scoccimarro et al., 2011)
CNRM-CM5	CNRM-CERFACS (France)	1.5 x 1.5	(Voldoire et al., 2012)
GFDL-ESM2M	NOAA-GFDL (USA)	2 x 2.5	(Dunne et al., 2012)
IPSL-CM5A-LR	IPSL (France)	1.875 x 3.75	(Dufresne & Bony, 2008)
MIROC5	MIROC (Japan)	2.8 x 2.8	(Watanabe et al., 2010)
MRI-CGCM3	MRI (Japan)	1.125 x 1.125	(Yukimoto et al., 2012)

We compute downscaled regional climate projections for the historical period of 1980–1999; from the RCP4.5 scenario, which assumes a moderate amount of mitigation of GHG emissions (van Vuuren et al., 2011) for two 20-year periods of the 21st century (2040–2059 and 2080–2099); and from the RCP8.5 scenario, which assumes only a minimum of mitigation, for the 2080–2099 period (Riahi et al., 2011). For the state of Minnesota and surrounding regions, we use WRF to generate hourly averages of the following surface variables: 2-m air temperature, 2-m humidity, 10-m wind speed and direction, precipitation, downward solar radiation, net radiation, latent, sensible, and ground heat fluxes, snow depth, and soil temperature and moisture content at 10 layers to a depth of 2.5 m. The seasonal cycle of these values, especially precipitation, is generally improved by physical downscaling (Mendez et al., 2020).

We perform a simple linear-scaling bias-adjustment (Teutschbein & Seibert, 2012) to surface air temperature and precipitation as described in equations 1–4 in Shrestha et al. (2017) using monthly mean observations as reported by the PRISM group (Daly et al., 2017). For snow depth, we perform bias adjustment with the NSIDC analysis that uses a combination of observed snow depth observations and the PRISM data set (Broxton et al., 2019; Zeng et al., 2018).

For the bias adjustment, we compare observations and WRF-simulated values from runs forced with each GCM for each monthly average (i.e., the average temperature difference over every January from 1980 to 1999 is calculated to receive one offset value for January at each grid point). Precipitation and snow depth are scaled by dividing monthly observational averages for 1981–2000 by monthly model values. The 2-m air temperature, precipitation, and snow depth error adjustments are then applied to data from each WRF run for each future scenario. Bias adjustment based on linear scaling retains the interannual variability but forces each multiyear monthly average for each GCM-forced WRF run in the historical simulations to equal the PRISM observations and NSIDC analysis, respectively. Linear scaling assumes that this offset carries through to the climate simulations of the future, so the simulations will now diverge in their calculations of these variables. Variables other than air temperature, precipitation, and snow depth are not adjusted because of lack of available observations. The prognostic variables for CCSM4 and CMCC-CM were previously bias-adjusted, so their historical multiyear monthly means match reanalysis data as described in Bruyère et al. (2014). We apply the same bias adjustment to the prognostic variables of all future scenarios for these two models.

In addition to analyzing WRF results forced from each GCM, we analyze the multimodel ensemble (MME) of each variable averaged over all WRF-driven



**Figure 1.** The outer (complete map) and inner grid (black frame) used for climate projections. The state of Minnesota is marked in white.



runs from all GCMs. Individual years of the simulations are treated as individual ensemble members in our analyses, and we adjusted the degrees of freedom in our statistical tests to account for lag-1 autocorrelation in our data, according to Wilks (2011), which allows a robust statistical analysis with 160 ensemble members per scenario. The advantage of this approach is that the variability of individual GCM simulations is being preserved, compared to smaller ensemble sizes with average GCM forcings. Although an MME approach with eight down-scaled models should be considered as more reliable than individual model results, as previously demonstrated by Pincus et al. (2008), we also quantify bias adjustments for individual realizations of the historical climate in the next section.

### 3. Results and Discussion

#### 3.1. Statewide Area Averages


The statewide area averages are computed by averaging over all grid cells with more than 50% of their area inside the state. Figure 2 depicts the statewide area averages for the bias adjustment offset for 2-m temperature and the adjustment factor for precipitation. These adjustments are applied to the WRF simulations to generate the bias-adjusted results. Although the GCM input data (Figures 2a and 2c) are closer to the observations, the WRF simulations (Figures 2b and 2d) convey the larger climate variability on the regional scale. However, the larger bias in the simulated precipitation is partly due to the choice of the 10-km horizontal resolution, which is within the 5–10 km range where neither convective parameterization nor a fully explicit approach provide convincing results (Molinari & Dudek, 1992). WRF-CLM4crop uses a convective parameterization scheme that adds subgrid-scale precipitation to the explicit approach and thus leads to an overestimation of precipitation in our results, in contrast to the study by Ashfaq et al. (2016), which uses cumulus convection parameterization on the larger 18 km grid or the study by Liu et al. (2017), which does not add cumulus convection parameterization to the smaller 4 km grid.

Although the trend analysis in the present paper is independent of linear scaling, we find that the analysis of absolute values and threshold values, such as the number of days per year with snow depth of more than a certain value benefits from the bias adjustment. It should also be noted that the downscaled results (Figures 2b and 2d) do not reflect the performance of individual GCMs (Figures 2a and 2c), they are merely a reflection of how WRF interprets the given sets of input data. Thus, although we use the term MME, our results are obtained with a single regional model at a single horizontal resolution, which makes the statistical analysis more feasible.

Many characteristics of individual GCM input data are also reflected in the WRF simulations during our 20-year historical simulation. For example, GFDL-ESM2M, IPSL-CM5A-LR, MIROC5, and MRI-CGCM3 have a warm bias in winter, whereas bcc-csm1-1 and CCSM4 have a cold bias in winter. CNRM-CM5 shows a slight warm bias from July to September, and MIROC5 depicts a strong warm bias from June to the end of the year. All WRF simulations apart from the ones driven by CMCC-CM have a cold bias in April and May, potentially due to the positive precipitation bias during this wet season. All area-averaged GCM input data are too dry in winter but only WRF runs driven by CMCC-CM, GFDL-ESM2M, IPSL-CM5A-LR, and MRI-CGCM3 share this GCM dry bias in winter. Especially in summer, all WRF runs are too wet based on the overrepresentation of convective precipitation, as discussed above (Figures 2c and 2d).

A detailed spatial analysis of the differences between the nonbias-adjusted WRF runs and the GCM input data in terms of comparability to observations is given in Figures S1 and S2 in the supplemental material. Figure S1 in Supporting Information S1 shows that historical 2-m temperatures in WRF are slightly too high in winter and too low in the spring rainy season compared with observation and also with GCM input. However, in summer, WRF simulations fit better with observations than the GCM input. Summer temperatures in the GCM input data are generally too high, but therefore match over the urban heat island of the Twin Cities metro area in southeast Minnesota, although urban climate is typically not included in GCMs. Fall temperatures are generally well represented in WRF and the GCM input data. Figure S2 in Supporting Information S1 depicts the general wet bias in the WRF MME that outweighs the aforementioned dry bias in some WRF simulations in winter. Although the relative precipitation bias is strongest in the dry summer season (Figure 2d), the absolute differences to observations are still not statistically significant over the southcentral region.

Figures S3–S6 in Supporting Information S1 compare future projections from the MMEs for GCMs and the WRF runs. Despite the wet bias in the WRF MME, they have similar trends for all three emission scenarios. Figure S7




**Figure 2.** Bias adjustments area-averaged over every grid cell within the state of Minnesota for (a) 2-m temperature difference between GCM runs and PRISM, (b) as (a) but for 2-m temperature difference between WRF runs and PRISM, (c) as (a) but for fraction of PRISM precipitation over GCM runs, and (d) as (b) but for fraction of PRISM precipitation over WRF runs. Please note that the optimum offset in (a) and (b) is 0, whereas the optimum fraction in (c) and (d) is 1.

in Supporting Information S1 provides the bias adjustment factors for snow depth. Their values divert away from 1.0 for smaller absolute snow depth values in spring and fall. We suggest that higher values in WRF snow depth are a direct result of increased precipitation.


MME 2-m temperature increases in each scenario compared to the historical period, particularly in winter (Figure 3). WRF simulates less future warming than the MME when driven by MRI-CGCM3 and GFDL-ESM2M





**Figure 3.** Monthly average 2-meter temperature averaged over every grid cell within the state of Minnesota for each Weather Research and Forecasting (WRF)-downscaled GCM (colors), the multimodel mean (MME; black line), and the PRISM data set (dashed line). Also shown are all years of the 20-year WRF simulations for all GCMs (160 realizations; gray lines).

and generally stronger warming when forced with MIROC5 and IPSL-CM5A-LR. Simulated warming driven by bcc-csm1-1, CCSM4, CMCC-CM, and CNRM-CM5 is relatively close to the MME. Figure S8 in Supporting Information S1 depicts the anomaly time series and overall standard deviation for a more detailed analysis of future trends, which does not only show the projected temperature increase of 2°C by the midcentury but also results in the individual WRF runs. These include strong warming during the month of May of up to 13°C in the



**Figure 4.** As Figure 3, but for precipitation.

MIROC5-driven runs and an especially weak warming during the month of March of generally less than  $1^{\circ}\text{C}$  in the MRI-CGCM3-driven runs.

Simulated precipitation variability (Figure 4) increases in spring and summer, especially in the late 21st century. Early summer MME rainfall increases in the mid-century and in the RCP8.5 late-century scenarios, while fall MME rainfall increases in both scenarios in the late century. There is a very small increase in winter MME precipitation of about  $0.1\text{ mm d}^{-1}$  in all scenarios. When WRF is driven with GFDL-ESM2M and MRI-CGCM3, increases in summer are strongest with values above  $2\text{ mm d}^{-1}$ , whereas the MIROC5-driven simulations show

decreases of up to  $2 \text{ mm d}^{-1}$  from late spring through early fall in all scenarios. WRF-forced simulations of precipitation from CMCC-CM are notably different from those of other models, with large increases of up to  $4 \text{ mm d}^{-1}$  in the fall for both late 21st-century simulations. As with temperature, precipitation from WRF driven with bcc-csm1-1, CCSM4, and CNRM-CM5 is closest to the MME. Anomaly time series and standard deviation for precipitation and the related snow depth are displayed in Figures S9 and S10 in Supporting Information S1.

### 3.2. Spatial Distributions


Historical 20-year average winter (Figure 5a) and summer (Figure 6a) MME 2-meter temperatures illustrate the strong north-south temperature gradient in the state. Anomalies of future winter MME projections (Figures 5b–5d) show that temperature increases are strongest along the northern border in the mid-century runs (Figure 5b), throughout much of the northern half of the state in RCP4.5 by the end of the century (Figure 5c) and throughout most of the northern half of the state in the RCP8.5 scenario (Figure 5d).

The increased rate of warming in the north is suggested to be related to both synoptic-scale warming (Wang et al., 2017; their Figure 8) as well as reduced albedo from reduced snow cover, which results in increases in average winter temperature ranging from  $\sim 1^\circ\text{C}$  by the mid-century to  $6^\circ\text{C}$  by the end of century in RCP8.5 (Figure 5d). This warming trend can be observed across Minnesota (Runkle et al., 2017; their Figure 1) and our simulations suggest that the trend will continue. Lakes will be ice-free for longer periods and the resulting decrease in albedo will contribute to local winter warming. The average summer temperature shows a more homogeneous increase across the state that ranges from  $\sim 1^\circ\text{C}$  by the mid-century (Figures 6b) to  $5^\circ\text{C}$  by the end of the century in the RCP8.5 scenario (Figure 6d). These simulated future increases in average summer temperature contrast with observations of average summer temperature across the state that do not have a significant trend in the historical record (Runkle et al., 2017; their Figure 2a).


Some of the future simulations also do not depict significant temperature changes by the mid-century, especially GFDL-ESM2M and MRI-CGCM3, whereas others show a much stronger warming. Strong mid-century warming of about  $4^\circ\text{C}$  occurs with bcc-csm1-1 and CNRM-CM5 forcing in winter and with IPSL-CM5A-LR and MIROC5 forcing in summer (Figures 3b and S8 in Supporting Information S1), which is also depicted in the detailed maps of temperature trends for different GCM input data (Figures S11 and S12 in Supporting Information S1). The stronger warming in the northern parts of the state in the MME is consistent with most model simulations, only CCSM4 and MRI-CGCM3 show a stronger warming in southern and central Minnesota during both seasons and GFDL-ESM2M only during summer. CNRM-CM5 and MIROC5 project a more pronounced negative west-east gradient in winter temperature change than most models, only simulations forced with MRI-CGCM3 and in RCP4.5 also CMCC-CM result in a positive west-east gradient during both seasons. In GFDL-ESM2M, this appears only in winter. Input data from CCSM4, CMCC-CM, and IPSL-CM5A-LR lead to the strongest response over lakes. In summer, increased evaporation over lakes mitigates local warming.

According to the statewide average analysis of precipitation (Figure 4), most of the change in future precipitation occurs in spring and early summer; therefore, we analyze spring average MME precipitation (rain and snow water equivalent) here. Spring average MME precipitation across Minnesota is strongest in the southeast portion of the state and weakest in the northwest (Figure 7a). Simulated precipitation changes by the mid-century differ among WRF runs with some runs showing spring increases (e.g., driven with IPSL-CM5A-LR) and some showing decreases (e.g., driven with MIROC5 and GFDL-ESM2M; Figure 4b), which together result in no significant changes in spring average precipitation across the state (Figure 7b). By the end of the century, spring precipitation is projected to increase slightly in the far north of the state in the RCP4.5 scenario (Figure 7c) and by up to  $1 \text{ mm d}^{-1}$  in the northern half of the state as well as the southern portion of the domain in Iowa in RCP8.5 (Figure 7d). This is in contrast to statistical projections from Localized Constructed Analogs (LOCA; Pierce et al., 2014), where the strongest precipitation increase occurs in the northeastern and central parts of the state (Figure S13 in Supporting Information S1). This is similar to the historical trend (NCEI, 2021). Our results show that the greatest future increase in rainfall is projected to occur in the northern part of the state indicate a change in regional circulation, and spring average rainfall across the state will become more homogeneous. Winter average MME precipitation is projected to increase slightly, with a statistically significant increase by end of century of up to  $0.25 \text{ mm d}^{-1}$  in RCP8.5 (not shown).







**Figure 5.** Average winter (December–February) MME 2-meter temperature in °C for (a) historical simulations and (b)–(d) anomalies of each RCP scenario compared to the historical period. Shading in (b)–(d) indicates statistically significant changes over U.S. land points at the 95% confidence interval. Please note that there is a different color bar for (a) than for (b)–(d).



**Figure 6.** As Figure 5, but for summer (June–August).



**Figure 7.** As Figure 5, but for precipitation and precipitation anomalies in  $\text{mm d}^{-1}$  in spring (March–May).



**Figure 8.** As Figure 5, but for MME snow depth and snow depth anomalies (cm) in winter (December–February).



Average winter MME snow depth generally increases with latitude across the state; however, there is a lobe of lower snow depths stretching northward on the far western side of the state (Figure 8a). Despite the currently observed and projected increases in precipitation, snow depth is projected to decrease across the state except in the northernmost region during the 21st century (Figures 8b–8d) because of increased surface air temperature. Strongest decreases in snow depth are projected to occur in central Minnesota, where average snow depth is expected to decrease by up to 50% by the middle of the 21st century (Figure 8b). By the end of the 21st century under RCP8.5, this change is expected to also cover southern Minnesota and the maximum reduction in snow depth reaches more than 12 cm. While snow depth over the northernmost part of the state remains virtually unchanged, the simulations show significant decreases in snow depth along the Minnesota North Shore and into much of Wisconsin exceeding 12 cm. These regions include part of the U.S. National Forest system and are at risk of decreasing revenues in winter recreation as well as threats to ecosystem health from pests that may survive warmer winters (Govindan & Hutchison, 2020; Venette & Hutchison, 2021).

The average MME number of days per year when snow depth meets or exceeds a threshold of 2.54 cm (equivalent to one inch) follows a similar pattern as average MME snow depth (Figure 9a). Warmer winters result in fewer days with snow cover on the ground. Significant decreases in days per year with snow cover above one inch are found over central and southeast Minnesota and western Wisconsin of up to 40 days per year by the mid-century (Figure 9b). By the end of the century in the RCP8.5 scenario, there are up to 55 fewer days of snow cover in Minnesota and more than 60 fewer days in central Wisconsin (Figure 9d). Because of the large uncertainty of observed snow depth, we also provide non adjusted snow depth values in Figures S14–S16 in Supporting Information S1. They complement Figures S10 in Supporting Information S1 and Figures 8 and 9, and although snow depth is generally lower in observations, the days per year with snow cover are remarkably similar, which suggests that the number of snow events is similar, despite the higher snowfall rate in the WRF simulations.

The goal of this study is to develop a dynamically downscaled climate data set for Minnesota to be used for impact studies. This data set will be useful if it provides value to previously developed, well-tested data sets (i.e., higher resolution while broadly agreeing with other projections). While there are differences in the number of GCMs, time periods, and downscaling methods between our study and that of the National Climate Assessment (NCA), our results are consistent in magnitudes and patterns. For example, our statewide average annual temperature increases (Figures 3b–3d) agree broadly with Midwest average projected increases for RCP4.5 by mid-century ( $\Delta 2.3^{\circ}\text{C}$ ) and for end of century for RCP4.5 ( $\Delta 3.1^{\circ}\text{C}$ ) and RCP8.5 ( $\Delta 5.3^{\circ}\text{C}$ ) in Table 6.4 of the NCA (Vose et al., 2017).


The benefits of downscaling are highlighted for a variable like precipitation that varies strongly in space and time and for which variability is projected to increase in the future. Like our analysis, the NCA found more significant changes in projected precipitation in winter and spring than in fall and summer (Easterling et al., 2017). While their analysis shows a homogeneous increase in winter and spring precipitation for RCP8.5 by the end of the century of about 20%, we found a smaller though statistically significant increase in winter precipitation and more spatial variability in projected spring precipitation. Our projected RCP8.5 end of the century spring precipitation ranges from no statistically significant change in the southern portion of the state to a 12%–30% increase in the central region and a 30%–60% increase in the north. While our analysis provides a more detailed projection of future precipitation than the NCA, we note that there are limitations to our analysis because of the single regional climate model used to downscale data and the single method of bias adjustment we employed (Laux et al., 2021). Future studies will examine alternate methods of bias adjustment of this data set.

The data presented here are immediately useful for impact studies of agricultural, energy, economic, and other ecosystem service sectors of Minnesota and will be a baseline for comparison with future downscaling efforts of CMIP6 (Stouffer et al., 2017). Multimodel ensemble values of temperature and precipitation are available at a variety of user-defined domains from [climate.umn.edu](http://climate.umn.edu) (The Minnesota Department of Natural Resources Climate Explorer Tool), and all daily data are available for download at the University of Minnesota Digital Conservancy website. Other raw data are available upon request.

#### 4. Conclusions

The present study describes a high-resolution regional climate modeling effort over the U.S. state of Minnesota that is already providing input for various projects, including improved projections of weather extremes, management of infrastructure, industry, and water resources (Noe et al., 2019) as well as the tracking of invasive species





**Figure 9.** As Figure 5, but for MME days per year and difference in days per year with snow depth above 2.54 cm (one inch).

(Govindan & Hutchison, 2020; Venette & Hutchison, 2021). This state-level downscaling effort links climate projections to decision-making within regional communities. Snow depth simulations emphasize the need for detailed modeling efforts of the hydrological cycle, especially over high-latitude climates.

## Data Availability Statement

The daily WRF model data are hosted at the University of Minnesota Digital Conservancy website at: <https://doi.org/10.13020/YV29-JY19>. CMIP5 forcing data were obtained from the World Data Center for Climate (WDCC) hosted by the German Climate Computing Center (DKRZ) at <https://cera-www.dkrz.de/WDCC/ui/cerasearch/q>. Monthly temperature and precipitation observations for 1981–2000 were made available by the PRISM group at <http://www.prism.oregonstate.edu/recent>, and the snow depth analysis was provided by NSIDC at <https://doi.org/10.5067/0GGPB220EX6A>.

## Acknowledgments

The authors acknowledge Keith J. Harding and the Minnesota Supercomputing Institute at the University of Minnesota for providing resources that contributed to the research results reported within this paper (<http://www.msi.umn.edu>). Funding for this project was provided by a grant from the Legislative-Citizen Commission on Minnesota Resources Grant Number: M. L. 2014, Chp. 312, Sec. 8 to the Minnesota Invasive Terrestrial Plants and Pests Center (<http://www.mitppc.umn.edu>), and by a grant from the Minnesota Environment and Natural Resources Trust Fund Grant Number: M. L. 2015, Chp. 76, Sec. 2, Subd. 04a (<http://www.legacy.mn.gov>) as recommended by the Legislative-Citizen Commission on Minnesota Resources. The Trust Fund is a permanent fund constitutionally established by the citizens of Minnesota to assist in the protection, conservation, preservation, and enhancement of the state's air, water, land, fish, wildlife, and other natural resources. Currently, 40% of net Minnesota State Lottery proceeds are dedicated to growing the Trust Fund and ensuring future benefits for Minnesota's environment and natural resources. The authors are grateful to three anonymous reviewers for their insightful comments.

## References

- Ashfaq, M., Rastogi, D., Mei, R., Kao, S.-C., Gangrade, S., Naz, B. S., & Touma, D. (2016). High-resolution ensemble projections of near-term regional climate over the continental United States. *Journal of Geophysical Research: Atmospheres*, 121, 9943–9963. <https://doi.org/10.1002/2016JD025285>
- Bekryaev, R. V., Polyakov, I. V., & Alexeev, V. A. (2010). Role of polar amplification in long-term surface air temperature variations and modern Arctic warming. *Journal of Climate*, 23, 3888–3906. <https://doi.org/10.1175/2010JCLI3297.1>
- Boé, J., Terray, L., Habets, F., & Martin, E. (2007). Statistical and dynamical downscaling of the Seine basin climate for hydro-meteorological studies. *International Journal of Climatology*, 27, 1643–1655. <https://doi.org/10.1002/joc.1602>
- Broxton, P., Zeng, X., & Dawson, N. (2019). Daily 4 km gridded SWE and snow depth from assimilated in-situ and modeled data over the conterminous US, version 1. NASA National Snow and Ice Data Center Distributed Active Archive Center. <https://doi.org/10.5067/0GGPB220EX6A>
- Bruyère, C. L., Done, J. M., Holland, G. J., & Fredrick, S. (2014). Bias corrections of global models for regional climate simulations of high-impact weather. *Climate Dynamics*, 43, 1847–1856. <https://doi.org/10.1007/s00382-013-2011-6>
- Callendar, G. S. (1938). The artificial production of carbon dioxide and its influence on temperature. *Quarterly Journal of the Royal Meteorological Society*, 64, 223–240. <https://doi.org/10.1002/qj.49706427503>
- Dai, Y., Zeng, X., Dickinson, R. E., Baker, I., Bonan, G. B., Bosilovich, M. G., et al. (2003). The common land model. *Bulletin of the American Meteorological Society*, 84, 1013–1023. <https://doi.org/10.1175/BAMS-84-8-1013>
- Daly, C., Slater, M. E., Roberti, J. A., Laseter, S. H., & Swift, L. W. (2017). High-resolution precipitation mapping in a mountainous watershed: Ground truth for evaluating uncertainty in a national precipitation dataset. *International Journal of Climatology*, 37, 124–137. <https://doi.org/10.1002/joc.4986>
- Dufresne, J.-L., & Bony, S. (2008). An assessment of the primary sources of spread of global warming estimates from coupled atmosphere–ocean models. *Journal of Climate*, 21, 5135–5144. <https://doi.org/10.1175/2008JCLI2239.1>
- Dunne, J. P., John, J. G., Adcroft, A. J., Griffies, S. M., Hallberg, R. W., Shevliakova, E., et al. (2012). GFDL's ESM2 global coupled climate–carbon Earth system models. Part I: Physical formulation and baseline simulation characteristics. *Journal of Climate*, 25, 6646–6665. <https://doi.org/10.1175/JCLI-D-11-00560.1>
- Easterling, D. R., Kunkel, K. E., & Arnold, J. R. (2017). Ch. 7: Precipitation change in the United States. *Climate Science Special Report: Fourth National Climate Assessment*, 1. <https://doi.org/10.7930/JOH993CC>
- Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., et al. (2011). The community climate system model version 4. *Journal of Climate*, 24, 4973–4991. <https://doi.org/10.1175/2011jcli4083.1>
- Giorgi, F., & Mearns, L. O. (1991). Approaches to the simulation of regional climate change: A review. *Review of Geophysics*, 29, 191. <https://doi.org/10.1029/90RG02636>
- Giorgi, F., Mearns, L. O., & Gutowski, W. J. (2015). Regional dynamical downscaling and the CORDEX initiative. *Annual Review of Environment and Resources*, 40, 467–490. <https://doi.org/10.1146/annurev-environ-102014-021217>
- Govindan, B. N., & Hutchison, W. D. (2020). Influence of temperature on age-stage, two-sex life tables for a Minnesota-acclimated population of the Brown marmorated stink bug (*Halyomorpha halys*). *Insects*, 11, 108. <https://doi.org/10.3390/insects11020>
- Harding, K. J., & Snyder, P. K. (2014). Examining future changes in the character of Central U.S. warm-season precipitation using dynamical downscaling. *Journal of Geophysical Research: Atmospheres*, 119(13), 13116–13136. <https://doi.org/10.1002/2014JD022575>
- Harding, K. J., Snyder, P. K., & Liess, S. (2013). Use of dynamical downscaling to improve the simulation of Central U.S. warm season precipitation in CMIP5 models. *Journal of Geophysical Research: Atmospheres*, 118(12), 12512–12536. <https://doi.org/10.1002/2013JD019994>
- Harding, K. J., Twine, T. E., VanLoocke, A., Bagley, J. E., & Hill, J. (2016). Impacts of second-generation biofuel feedstock production in the central U.S. on the hydrologic cycle and global warming mitigation potential. *Geophysical Research Letters*, 43(10), 10773–10781. <https://doi.org/10.1002/2016GL069981>
- Harvey, B. J., Shaffrey, L. C., & Woollings, T. J. (2015). Deconstructing the climate change response of the Northern Hemisphere wintertime storm tracks. *Climate Dynamics*, 45, 2847–2860. <https://doi.org/10.1007/s00382-015-2510-8/FIGURES/8>
- IPCC. (2013). *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*. T. F. Stocker, et al., Eds. (p. 1535). Cambridge University Press.
- IPCC. (2021). *Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*. In Masson-Delmotte, V., et al. (Eds.), (p. 3949). Cambridge University Press. Retrieved from <https://www.ipcc.ch/report/ar6/wg1/>
- Laux, P., Rötter, R. P., Webber, H., Dieng, D., Rahimi, J., Wei, J., et al. (2021). To bias correct or not to bias correct? An agricultural impact modelers' perspective on regional climate model data. *Agricultural and Forest Meteorology*, 304–305, 108406. <https://doi.org/10.1016/j.agrformet.2021>
- Liu, C., Ikeda, K., Rasmussen, R., Barlage, M., Newman, A. J., Prein, A. F. (2017). Continental-scale convection-permitting modeling of the current and future climate of North America. *Climate Dynamics*, 49(1/49), 71–95. <https://doi.org/10.1007/s00382-016-3327-9>

- Lu, Y., Jin, J., & Kueppers, L. M. (2015). Crop growth and irrigation interact to influence surface fluxes in a regional climate - Cropland model (WRF3.3—CLM4crop). *Climate Dynamics*, 45, 3347–3363. <https://doi.org/10.1007/s00382-015-2543-z>
- McGinnis, S., & Mearns, L. (2021). Building a climate service for North America based on the NA-CORDEX data archive. *Climate Services*, 22, 100233. <https://doi.org/10.1016/j.cliser.2021.100233>
- Mearns, L. O., Gutowski, W., Jones, R., Leung, R., McGinnis, S., Nunes, A., & Qian, Y. (2009). *A regional climate change assessment Program for north America*. Eos, Transactions American Geophysical Union, 90, 311. <https://doi.org/10.1029/2009EO360002>
- Mendez, M., Maathuis, B., Hein-Griggs, D., & Alvarado-Gamboa, L.-F. (2020). Performance evaluation of bias correction methods for climate change monthly precipitation projections over Costa Rica. *Water*, 12, 482. <https://doi.org/10.3390/w12020>
- Molinari, J., & Dudek, M. (1992). Parameterization of convective precipitation in mesoscale numerical models: A critical review. *Monthly Weather Review*, 120, 3262–3344. [https://doi.org/10.1175/15200493\(1992\)120<0326:pocpim>2.0.co;2](https://doi.org/10.1175/15200493(1992)120<0326:pocpim>2.0.co;2)
- NCEI. (2021). *National trends temperature, precipitation, and drought*. National Centers for Environmental Information (NCEI). Retrieved from <https://www.ncdc.noaa.gov/temp-and-precip/us-trends/tavg/win>
- Noe, R., Keeler, B., Twine, T., Brauman, K., Mayer, T., & Rogers, M. (2019). *Climate change projections for improved management of infrastructure, industry, and water resources in Minnesota*. Retrieved from <http://hdl.handle.net/11299/209130>
- Pierce, D. W., Cayan, D. R., & Thrasher, B. L. (2014). Statistical downscaling using localized constructed Analogs (LOCA). *Journal of Hydro-meteorology*, 15, 2558–2585. <https://doi.org/10.1175/JHM-D-14-0082.1>
- Pincus, R., Batstone, C. P., Hofmann, R. J. P., Taylor, K. E., & Glecker, P. J. (2008). Evaluating the present-day simulation of clouds, precipitation, and radiation in climate models. *Journal of Geophysical Research*, 113, D14209. <https://doi.org/10.1029/2007jd009334>
- Polyakov, I. V., Alekseev, G. V., Bekryaev, R. V., Bhatt, U., Colony, R. L., Johnson, M. A., et al. (2002). Observationally based assessment of polar amplification of global warming. *Geophysical Research Letters*, 29, 1878. <https://doi.org/10.1029/2001GL011111>
- Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., et al. (2011). RCP 8.5—A scenario of comparatively high greenhouse gas emissions. *Climate Change*, 109, 33–57. <https://doi.org/10.1007/s10584-011-0149-y>
- Runkle, J., Kunkel, K., Frankson, R., Easterling, D., & Champion, S. (2017). *Minnesota state climate summary*. NOAA Technical Report NESDIS 149-MN. 4. pp Retrieved from <https://statesummaries.ncics.org/chapter/mn/>
- Scoccimarro, E., Gualdi, S., Bellucci, A., Sanna, A., Giuseppe Fogli, P., Manzini, E. (2011). Effects of tropical cyclones on ocean heat transport in a high-resolution coupled general circulation model. *Journal of Climate*, 24, 4368–4384. <https://doi.org/10.1175/2011JCLI4104.1>
- Shi, X., Déry, S. J., Groisman, P. Y., & Lettenmaier, D. P. (2013). Relationships between recent pan-Arctic snow cover and hydroclimate trends. *Journal of Climate*, 26, 2048–2064. <https://doi.org/10.1175/JCLI-D-12-00044.1>
- Shrestha, M., Acharya, S. C., & Shrestha, P. K. (2017). Bias correction of climate models for hydrological modelling – Are simple methods still useful? *Meteorological Applications*, 24, 531–539. <https://doi.org/10.1002/met.1655>
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., et al. (2008). *A description of the Advanced Research WRF version 3* (p. 125). National Center for Atmospheric Research. <http://doi.org/10.5065/D68S4MVH>
- Stouffer, R. J., Eyring, V., Meehl, G. A., Bony, S., Senior, C., Stevens, B., & Taylor, K. E. (2017). CMIP5 scientific gaps and recommendations for CMIP6. *Bulletin of the American Meteorological Society*, 98, 95–105. <https://doi.org/10.1175/BAMS-D-15-00013.1>
- Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, 93, 485–498. <https://doi.org/10.1175/BAMS-D-11-00094.1>
- Teutschbein, C., & Seibert, J. (2012). Bias correction of regional climate model simulations for hydrological climate-change impact studies: Review and evaluation of different methods. *Journal of Hydrology*, 456–457, 12–29. <https://doi.org/10.1016/j.jhydrol.2012.05.052>
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., et al. (2011). The representative concentration pathways: An overview. *Climate Change*, 109, 5–31. <https://doi.org/10.1007/s10584-011-0148-z>
- Venette, R. C., & Hutchison, W. D. (2021). Invasive insect species: Global challenges, strategies & opportunities. *Frontiers Insect Science*, 1, 650520. <https://doi.org/10.3389/finsc.2021.650520>
- Voldoire, A., Sanchez-Gomez, E., Salas y Mélia, D., Decharme, B., Cassou, C., Séné, S., et al. (2012). The CNRM-CM5.1 global climate model: Description and basic evaluation. *Climate Dynamics*, 1–31. <https://doi.org/10.1007/s00382-011-1259-y>
- Vose, R. S., Easterling, D. R., Kunkel, K. E., LeGrande, A. N., & Wehner, M. F. (2017). Chapter 6: Temperature changes in the United States. *Climate Science Special Report: Fourth National Climate Assessment* (Vol. I). <https://science.2017.globalchange.gov/chapter/6/>
- Wang, J., Kim, H. M., & Chang, E. K. M. (2017). Changes in northern hemisphere winter storm tracks under the background of Arctic amplification. *Journal of Climate*, 30, 3705–3724. <https://doi.org/10.1175/JCLI-D-16-0650.1>
- Watanabe, M., Suzuki, T., Oishi, R., Komuro, Y., Watanabe, S., Emori, S., et al. (2010). Improved climate simulation by MIROC5: Mean states, variability, and climate sensitivity. *Journal of Climate*, 23, 6312–6335. <https://doi.org/10.1175/2010jcli3679.1>
- Wilks, D. S. (2011). In R. Dmowska, D. Hartmann, & H. T. Rossby (Eds.), *Statistical methods in the atmospheric sciences* (3rd ed., p. 676). Academic Press.
- Williams, D. N., Ananthakrishnan, R., Bernholdt, D. E., Bharathi, S., Brown, D., Chen, M., et al. (2009). The earth system grid: Enabling access to multimodel climate simulation data. *Bulletin of the American Meteorological Society*, 90, 195–205. <https://doi.org/10.1175/2008BAMS2459.1>
- Wu, T., Yu, R., Zhang, F., Wang, Z., Dong, M., Wang, L., et al. (2010). The Beijing climate center atmospheric general circulation model: Description and its performance for the present-day climate. *Climate Dynamics*, 34, 123–147. <https://doi.org/10.1007/s00382-008-0487-2>
- Xu, Z., Han, Y., & Yang, Z. (2018). Dynamical downscaling of regional climate: A review of methods and limitations. *Science China Earth Sciences*, 62, 62365–62375. <https://doi.org/10.1007/S11430-018-9261-5>
- Yukimoto, S., Adachi, Y., Hosaka, M., Sakami, T., Yoshimura, H., Hirabara, M., et al. (2012). A new global climate model of the meteorological research Institute: MRI-CGCM3—Model description and basic performance—. *Journal of Meteorological Society*, 90A, 23–64. <https://doi.org/10.2151/jmsj.2012-A02>
- Zeng, X., Broxton, P., & Dawson, N. (2018). Snowpack change from 1982 to 2016 over conterminous United States. *Geophysical Research Letters*, 45(12), 12940–12947. <https://doi.org/10.1029/2018GL079621>
- Zhang, X., Friedl, M. A., & Schaaf, C. B. (2006). Global vegetation phenology from Moderate Resolution Imaging Spectroradiometer (MODIS): Evaluation of global patterns and comparison with in situ measurements. *Journal of Geophysical Research*, 111, G04017. <https://doi.org/10.1029/2006jg000217>
- Zhou, W., Leung, L. R., Song, F., & Lu, J. (2021). Future changes in the Great Plains low-level Jet governed by seasonally dependent pattern changes in the north Atlantic subtropical high. *Geophysical Research Letters*, 48, e2020GL090356. <https://doi.org/10.1029/2020GL090356>
- Zorita, E., & von Storch, H. (1999). The analog method as a simple statistical downscaling technique: Comparison with more complicated methods. *Journal of Climate*, 12, 2474–2489. [https://doi.org/10.1175/15200442\(1999\)012<2474:tamaas>2.0.co;2](https://doi.org/10.1175/15200442(1999)012<2474:tamaas>2.0.co;2)